

## **Pap Smear Guidelines: Family Planning Update 2008 Part Two**

Produced by the Alabama Department of Public Health  
Video Communications and Distance Learning Division

## **Faculty**

Seshu P. Sarma, MD, FAAP  
Emory University Regional  
Training Center  
Atlanta, Georgia

## **Objectives**

- Discuss pap smear reporting
- Describe the Bethesda system 2006
- Discuss management of unsatisfactory smears and smears with no T zone component

## **Objectives**

- Discuss management strategies for various squamous and glandular cell abnormalities
- Discuss treatment modalities for CIN

## **Cervical Cancer**

- Pap smear screening resulted in dramatic reduction in cervical cancer
- The value of accurate screening can be reduced by loss to follow-up or under-treatment of significant lesions

## **Pap Smear Reporting**

- Pap system of reporting
  - Bethesda 1988 system
  - Bethesda 1991 system
  - Bethesda 2001 system
  - Bethesda 2006 system

### **Pap Reporting**

- Class 1
  - Normal
- Class 2
  - Atypia, inflammation, reactive change
- Class 3
  - Dysplasia
    - Mild, moderate and severe
- Class 4
  - Carcinoma in situ
- Class 5
  - Invasive cancer

### **WHO Reporting**

- CIN 1
  - Mild dysplasia
- CIN 2
  - Moderate dysplasia
- CIN 3
  - Severe dysplasia
  - Carcinoma in situ

### **Bethesda System**

- Bethesda 1988
- Bethesda 2001
- Bethesda 2006

### **Bethesda 2006**

- 146 experts
- 29 organizations
- Met September 18-19, 2006 in Bethesda, MD
- Developed revised, evidence based consensus guidelines for management of abnormal pap smears

### **Bethesda 2006**

- ASCUS
- LGSIL
  - Management unchanged for the above 2 categories except for adolescents
  - Cytologic follow-up for 2 years approved for adolescents

### **Bethesda 2006**

- Recommendations for HSIL and AGC underwent only minor changes
- More emphasis placed on immediate screen and treat for HSIL lesions

### **Bethesda 2006**

- HPV testing was incorporated into the management of AGC after initial evaluation with colposcopy and endometrial sampling

### **Bethesda 2006**

- The 2004 interim guidance for HPV testing as a adjunct to cervical cytology for screening in women 30 years of age or older was formally approved

### **Adequate Specimen**

- Conventional Smear:
  - 8,000-12,000 well visualized cells should be seen
- Liquid-based Pap:
  - 5,000 well visualized cells should be seen
  - Epithelial cells may be obscured by inflammation
  - If >75% of cells are obscured, it is considered unsatisfactory

### **Adequacy of Specimen**

- Satisfactory specimen contains
  - 10 well preserved endocervical cells
  - or
  - 10 squamous metaplastic cells

### **Transformation Zone Not Present**

- If Pap is otherwise normal, no higher risk of subsequent detection of high grade SIL
- Most women without an endocervical or transformation zone component can be re-screened in 12 months

### **No T Zone Components**

- Repeat testing is recommended for women at higher risk of neoplasia such as:
  - Immunocompromised
  - HPV +
  - previous abnormal pap smears
  - Inability to visualize the cervix
  - Insufficient previous screening

### **Unsatisfactory Pap Smears**

- Pap smears that are unsatisfactory need to be repeated in 2-4 months
- If three consecutive specimens are unsatisfactory, colposcopy is recommended
  - Histologic abnormalities were found in 16% of this population

### **Abnormal Pap Smears**

- 50-60 million pap tests per year in US
- 3.5 million are classified as abnormal and require some form of medical follow-up
- 10,000 new cases of cervical cancer seen each year in the US
- 4,000 deaths from cervical cancer

### **Abnormal Pap and Cervical Cancer**

- Determining which women with abnormal Pap tests are at risk of significant cervical disease and treating them presents a:
  - major public health challenge
  - multibillion dollar cost to our healthcare system

### **Inconclusive Pap Smears**

- Almost 2 million Pap smears each year are classified as “inconclusive”
  - ASC-US, ASC-H and LSIL
- Most of them have no detectable problem
- Less than one fifth of these women have a significant precancerous lesion

### **Inconclusive Pap Smears**

- Although these changes are mild, they may result in:
  - Anxiety
  - Unnecessary medical procedures
  - Extra costs to healthcare system

### **Abnormal Pap Smears**

- Effective cervical cancer prevention requires
  - Recognition and treatment of the precursors of invasive cancer
- The 2001 Bethesda System of nomenclature describes the categories of epithelial cell abnormalities seen on pap smears

### **2001 Bethesda System**

- Squamous Cell:
  - Atypical Squamous Cells
    - ASC-US (undetermined significance)
    - ASC-H ( can not exclude HSIL)
  - Low-grade squamous intraepithelial lesions
    - Mild dysplasia
    - HPV changes

### **2001 Bethesda System**

- Squamous Cell:
  - High-grade squamous intraepithelial lesions
    - Moderate and severe dysplasia and carcinoma-in-situ (CIN II and CIN III)
  - Squamous cell carcinoma

### **2001 Bethesda System**

- Glandular Cell:
  - Atypical glandular cells (AGC)
    - Endocervical
    - Endometrial
    - Not otherwise specified (NOS)

### **2001 Bethesda System**

- Glandular Cell:
  - Atypical glandular cells
    - Favor neoplasia
      - Specify endocervical
      - Not otherwise specified (NOS)
  - Endocervical adenocarcinoma in situ (AIS)
  - Adenocarcinoma

### **Precancerous Precursors**

- CIN 2, CIN 3, adeno-carcinoma in situ (AIS) are collectively referred to as CIN2/3+
- Cancer precursors include:
  - CIN 3, AIS and to a lesser extent CIN 2

### **HPV and Abnormal Paps**

- The presence of HPV is a marker for the risk of diagnosis of CIN2/3+
- Only 1 in 10 to 1 in 30 HPV infections are associated with abnormal cervical cytology
- Even a smaller proportion are associated with CIN2/3+

## HPV

- Among women with negative cytology and a positive HPV test result, only 15% will have abnormal cytology results within 5 years

## HPV

- HPV can express as CIN within months after infection
- However, the time course from CIN 3 to invasive cancer averages between 8.1 and 12.6 years

## CIN Regression

- Regression for CIN
  - 60% for CIN 1 and 40% for CIN 2

## CIN 1

- CIN 1 is the histologic appearance of cells producing HPV
- The goals of cervical cancer screening:
  - Not to prevent CIN
  - But to prevent and treat
    - Early invasive cervical cancer
    - Reduce mortality

## Management of Abnormal Pap

- Atypical Squamous Cells:
- The risk of cancer and CIN 2/3+ is low
  - 0.1-0.2% for cancer
  - 6.4-11.9% for CIN2/3+

## Atypical Squamous Cells

- Options include:
  - Immediate colposcopy
    - Provides rapid diagnosis but expensive
  - HPV DNA testing triage
  - Repeat cytology at 6 and 12 months

### **HPV Testing**

- Hybrid Capture DNA 2 assay for 13 oncogenic HPBV sub types
- Performed on the liquid based pap specimen
- Usually done within 3 weeks of obtaining the specimen

### **ASCUS (HPV+) in Adolescents**

- The risk of invasive cancer approaches zero in adolescents
- HPV infections are common and self limiting in adolescents
- Colposcopic evaluation is not necessary
- Can be monitored with pap smears at 6 and 12 months

### **Adolescents**

- Recommendations are similar for ASC and LSIL in adolescent girls
- LSIL represents HPV infection and most adolescents clear HPV and LSIL
- Follow-up Pap at 6 and 12 months is adequate

### **ASCUS and LSIL in Adolescents**

- Follow-up with cytology or HPV for up to 2 years is reasonable according to Bethesda 2006 guidelines

### **LSIL**

- More frequent in younger populations with larger number of recent partners
- Represents the appearance of cells that are actively engaged in HPV replication
- HPV testing is of limited value

### **LSIL**

- The risk of CIN 2/3+ at initial colposcopy following an LSIL is between 15% and 30%
- Colposcopy is recommended for these women
- For adolescents with LSIL, follow-up with Pap at 6 and 12 months is reasonable

### **LSIL in Adolescents**

- Follow-up at 6 and 12 months
  - Those with HSIL need colposcopy
- Follow-up at 24 months
  - Those with ASCUS or greater need colposcopy
    - Bethesda 2006 guidelines

### **LSIL in Postmenopausal Women**

- Options include:
  - Reflex HPV DNA testing
  - Repeat cytology at 6 and 12 months
  - Colposcopy
    - 2006 Bethesda guidelines

### **ASC-H**

- Atypical squamous cells-cannot exclude HSIL (ASC-H);
  - Includes 5-10% of ASC Pap smears
  - CIN2/3+ seen in 24-94% of such cases
  - Immediate colposcopic intervention is necessary
  - HPV testing is not cost effective

### **HSIL**

- Among women with HSIL cytology results:
  - CIN 2 and CIN 3 have been reported in 70% or more cases
  - Invasive cancer is seen in 1-2% of cases
  - Colposcopy with endocervical evaluation should be done
  - Entire vagina should be examined
  - See and treat approach can be followed if lesion seen

### **AGC**

- Atypical Glandular Cells:
  - Histological correlation:
    - 9-38% of women with AGC have significant neoplasia
    - 3-17% have cancer
    - CIN is the most common lesion associated with AGC

### **AGC**

- Source of the lesion
  - Cervix
  - Endocervix
  - Endometrium



### **AGC**

- Evaluation includes:
  - Colposcopy
  - Endocervical curettage
  - Endometrial sampling
    - With abnormal endometrial cells
    - Women >35 years of age
    - Younger women with abnormal bleeding, morbid obesity and oligomenorrhea

### **AGC**

- When evaluation is negative and histology is negative, follow-up includes:
  - AGC favor neoplasia: excision is warranted
    - Cold knife cone is a good choice
  - AGC and AGC NOS
    - Pap and ECC every 6 months until 4 consecutive negative smears are obtained

### **How Should CIN 1 Be Managed?**

- Depends on the preferences of the
  - Physician and the patient
  - For most young women, observation seems appropriate
  - Follow-up with pap smear every 6 months X 2
    - Colposcopy for an ASC or higher
    - or a positive HPV at 12 months

### **CIN 2 and 3**

- CIN 2 and 3 are recognized potential cancer precursors
  - Although CIN2 is associated with significant spontaneous regression
  - 40% of CIN 2 cases regress over 2 years
- Immediate treatment of CIN2 and CIN 3 with excision or ablation is recommended in non-pregnant patients

### **CIN 2 in Adolescents**

- Care of the adolescent with CIN 2 may be individualized
- Treatment may be deferred
- Close follow-up with pap and colposcopy may be adequate

### **Management of AIS**

- Cold-knife conization is recommended
  - To preserve specimen orientation
  - Permit optimal interpretation of histology and margin status
- LEEP is not recommended

### **Hysterectomy for CIN 2/3+**

- Hysterectomy is not the initial treatment of choice for
  - CIN 2 or 3
  - May be considered for persistent or recurrent CIN 2 or 3

### **Abnormal Pap in Pregnancy**

- ASC or LSIL:
  - Colposcopy during pregnancy or at 6-12 weeks postpartum
- ASC-H
  - Immediate colposcopy

### **Abnormal Pap in Pregnancy**

- HSIL
  - Immediate colposcopy
- AGC
  - Immediate colposcopy
  - No endocervical sampling is done during pregnancy

### **Pregnancy**

- CIN is not treated in pregnancy
- Excision should be considered
  - If lesion is suspicious for invasive CA
- Repeat colposcopic evaluation is done during pregnancy as needed
- Reassessment by colposcopy
  - 6-12 weeks postpartum

### **Treatment Modalities for CIN**

- Cryotherapy
- Laser ablation
- LEEP
- Cold knife cone biopsy
  - Cryotherapy or laser ablation are done only when endocervical pathology is ruled out

### **See and Treat Approach**

- During colposcopic evaluation, when significant abnormality is found, LEEP is performed to both evaluate and treat the precancerous condition

## Management of Cervical Disease

- Diagnostic Procedures:
  - HPV DNA testing
  - Colposcopy
  - LEEP (both diagnostic and therapeutic)
  - Cone biopsy (both diagnostic and therapeutic)

## Hybrid Capture DNA Assay

- Hybrid Capture DNA 2 assay
- Designed to detect 13 high-risk HPV subtypes (16,18,31,33,35,39,45,51,52,56,58,59,68)

## HPV DNA (High Risk) Testing

- Potential indications:
  - Triage of patients with ASCUS
  - Surveillance of HSIL
  - Combination Pap/DNA testing

## Self Collected HPV DNA

- Studies suggest that self-collected specimens for HPV testing may offer
  - A means to increase screening
  - Primary screening of older women

## Fournier Self Collection Device



## Colposcopy

- Evaluation of the
  - Cervix and vagina
  - Vulva
- With 10 – 25 times magnification
- Acetic acid 3-6 % solution used
- Cervix and vagina are evaluated before and after acetic acid is applied

## Colposcopy

- Pap smear is a screening test
- Colposcopy is the diagnostic test to confirm the disease process



## Colposcopy

- Colposcopy is performed when:
  - The pap smear is abnormal
  - The cervix looks abnormal
  - Women are exposed to DES in utero

## Treatment

- Cryotherapy
- Laser
- Cone biopsy
- LEEP (Loop Electrical Excision Procedure)

## Treatment of Cervical Disease

- Treatment of the precancerous disease depends on the
  - Severity of the disease
    - Mild, moderate or severe dysplasia or CIS
  - Involvement of the endocervical canal
  - On the age of the patient such as an adolescent

## Cryotherapy

- Destruction of abnormal tissue by freezing
- Cryotherapy destroys normal tissue along with the abnormal tissue

### **Cryosurgery**



### **Cryo Tip**



### **Cone Biopsy**

- Cone biopsy is an extensive form of cervical biopsy
- A wedge shaped tissue is removed from the cervix and examined under a microscope

### **Cone Biopsy**

- Cone biopsy removes abnormal tissue that is high in the cervical canal
- A small amount of normal tissue around the abnormal tissue is removed so that a margin free of abnormal cell is left in the cervix

### **Cold Knife Cone Biopsy**



### **Cold Knife Cone Biopsy**



### Cold Knife Cone Biopsy



### Cold Knife Cone Biopsy




### Cone Biopsy

- Cone biopsy can be done using
  - A surgical knife (cold knife cone)
  - A carbon dioxide laser
  - LEEP (Loop Electrical Excision Procedure)


### LEEP/LLETZ

- Loop Electro-surgical Excision Procedure, also know as large loop excision of the transformation zone is a technique used for treatment and diagnosis of cervical dysplasia.

### LEEP



### LEEP



## LEEP



## LEEP



## LEEP

- **LEEP**
  - Performed as an outpatient technique at the gynecologist's office except under special circumstances
  - The cervix is numbed, and a portion of the cervix is removed with a thin wire loop carrying a small electrical current
  - The tissue is then sent for pathologic evaluation

## Counseling

- Emphasize practicing safer sex in the future
- Encourage involvement of partners in the discussion
- Provide culturally sensitive and appropriate educational materials that are helpful
- Give them opportunity to return and discuss the information at a later time

## Cervical Cancer

- HPV is the primary cause of cervical cancer
- In most cases, the HPV is harmless and causes no symptoms
- The majority of women with HPV will not develop cervical cancer
- Cervical cancer is completely preventable if precancerous changes are detected and removed from the cervix

## References

- The HPV DNA virus hybrid capture assay:
- What is it-and where do we go from here? [www.mlo-online.com](http://www.mlo-online.com) (March 2003)
- Cervical cancer: prevention, diagnosis, and therapeutics:[www.cancer.org](http://www.cancer.org)
- Human Papillomaviruses: applications, caveats and prevention: The journal of Reproductive Medicine: Volume 47, No. 7/July 2002

## **References**

- **ACOG Practice Bulletin: Clinical Management Guidelines for Obstetrician-Gynecologists Number 66, September 2005**
- **Bethesda 2006 guidelines**