

Ordering Information for Rabies Immune Globulin (RIG) and Rabies Vaccine

Rabies Immune Globulin (RIG) and rabies vaccine can be ordered by a licensed physician or through a pharmacist directly from the manufacturer. At this time, ADPH is recommending the use of “Imovax” from Sanofi Pasteur or “RabAvert” from Novartis. Use of a licensed Rabies Immune Globulin is also available from each manufacturer.

Sanofi Pasteur 1-800-VACCINE

Novartis 1-877-683-4732

Programs for Uninsured and Underinsured Patients

Patient assistance programs that provide medications to uninsured or underinsured patients are available for rabies vaccine and Immune globulin.

Sanofi Pasteur's Patient Assistance Program (providing Imogam ® Rabies-HT and Imovax ® Rabies as well as other vaccines) is now administered through the Franklin Group. A healthcare professional or patient can either contact the Franklin Group directly, or call the customer service team (1-800-VACCINE) who will transfer them to the Franklin Group. The Franklin Group will review the application against the eligibility criteria. For more information about the program or to request an application, please contact the Sanofi Pasteur, Inc. Patient Assistance Program (Franklin Group) at 1 (866) 801-5655.

Novartis' Patient Assistance Program for RabAvert ® is managed through RX for Hope and can be accessed at 1-800-589-0837. Instructions and request forms are also available at the Rx for Hope website RabAvert Patient Assistance Program. Instructions and request forms are also available at the Sanofi Patient Connection website.

Variation from Human Rabies Vaccine Package Inserts

These new ACIP 5-dose series for PEP recommendations differ from current rabies vaccine label instructions. Historically, ACIP review and subsequent public health recommendations for the use of various biologics have occurred after vaccine licensure and generally are in agreement with product labels. However, differences between ACIP recommendations and product labels are not unprecedented. For example, during the early 1980s, ACIP review and recommendations concerning the intradermal use of rabies vaccines occurred well in advance of actual label claims and licensing (9). Based on discussions with industry representatives, alterations of current product labels for HDCV and PCEC are not anticipated by the producers of human rabies vaccines licensed for use in the United States.

TABLE 2: Rabies postexposure prophylaxis (PEP) schedule --- United States, 2010

Vaccination status	Intervention	Regimen*
Not previously vaccinated	Wound cleansing	All PEP should begin with immediate thorough cleansing of all wounds with soap and water. If available, a virucidal agent (e.g., povidine-iodine solution) should be used to irrigate the wounds.
	Human rabies immune globulin (HRIG)	Administer 20 IU/kg body weight. If anatomically feasible, the full dose should be infiltrated around and into the wound(s), and any remaining volume should be administered at an anatomical site (intramuscular [IM]) distant from vaccine administration. Also, HRIG should not be administered in the same syringe as vaccine. Because RIG might partially suppress active production of rabies virus antibody, no more than the recommended dose should be administered.
	Vaccine	Human diploid cell vaccine (HDCV) or purified chick embryo cell vaccine (PCECV) 1.0 mL, IM (deltoid area†), 1 each on days 0,§ 3, 7 and 14.¶
Previously vaccinated**	Wound cleansing	All PEP should begin with immediate thorough cleansing of all wounds with soap and water. If available, a virucidal agent such as povidine-iodine solution should be used to irrigate the wounds.
	HRIG	HRIG should not be administered.
	Vaccine	HDCV or PCECV 1.0 mL, IM (deltoid area†), 1 each on days 0§ and 3.

* These regimens are applicable for persons in all age groups, including children.

† The deltoid area is the only acceptable site of vaccination for adults and older children. For younger children, the outer aspect of the thigh may be used. Vaccine should never be administered in the gluteal area.

§ Day 0 is the day dose 1 of vaccine is administered.

¶ For persons with immunosuppression, rabies PEP should be administered using all 5 doses of vaccine on days 0, 3, 7, 14, and 28.

** Any person with a history of pre-exposure vaccination with HDCV, PCECV, or rabies vaccine adsorbed (RVA); prior PEP with HDCV, PCECV or RVA; or previous vaccination with any other type of rabies vaccine and a documented history of antibody response to the prior vaccination.